



Draft Genome Sequence of *Shewanella* sp. Strain P1-14-1, a Bacterial Inducer of Settlement and Morphogenesis in Larvae of the Marine Hydroid *Hydractinia echinata*

Citation

Rischer, Maja, Jonathan L. Klassen, Thomas Wolf, Huijuan Guo, Ekaterina Shelest, Jon Clardy, and Christine Beemelmans. 2016. "Draft Genome Sequence of *Shewanella* sp. Strain P1-14-1, a Bacterial Inducer of Settlement and Morphogenesis in Larvae of the Marine Hydroid *Hydractinia echinata*." *Genome Announcements* 4 (1): e00003-16. doi:10.1128/genomeA.00003-16. <http://dx.doi.org/10.1128/genomeA.00003-16>.

Published Version

doi:10.1128/genomeA.00003-16

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:25658465>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Draft Genome Sequence of *Shewanella* sp. Strain P1-14-1, a Bacterial Inducer of Settlement and Morphogenesis in Larvae of the Marine Hydroid *Hydractinia echinata*

Maja Rischer,^a  Jonathan L. Klassen,^b Thomas Wolf,^a Huijuan Guo,^a Ekaterina Shelest,^a Jon Clardy,^c Christine Beemelmans^a

Leibniz Institute for Natural Product Research and Infection Biology e.V., Jena, Germany^a; Department of Molecular & Cell Biology, University of Connecticut, Storrs, Connecticut, USA^b; Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, Massachusetts, USA^c

M.R. and J.L.K. contributed equally to this article.

The assembly and annotation of the draft genome sequence of *Shewanella* sp. strain P1-14-1 are reported here to investigate the genes responsible for interkingdom interactions, secondary metabolite production, and microbial electrogenesis.

Received 3 January 2016 Accepted 5 January 2016 Published 18 February 2016

Citation Rischer M, Klassen JL, Wolf T, Guo H, Shelest E, Clardy J, Beemelmans C. 2016. Draft genome sequence of *Shewanella* sp. strain P1-14-1, a bacterial inducer of settlement and morphogenesis in larvae of the marine hydroid *Hydractinia echinata*. Genome Announc 4(1):e00003-16. doi:10.1128/genomeA.00003-16.

Copyright © 2016 Rischer et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Christine Beemelmans, christine.beemelmans@hki-jena.de.

Members of the genus *Shewanella* are facultative anaerobic bacteria colonizing mostly marine and freshwater environments (1, 2) and have been shown to be metal-reducing bacteria capable of transferring electrons to the environment by different mechanisms (3). Besides their bioremediation capabilities and electrogenic potential, *Shewanella* spp. are known to produce secondary metabolites and induce settlement in larvae of biofouling marine invertebrates (4, 5). In our quest to isolate and characterize chemical cues involved in the larval settlement of *Hydractinia echinata* (6) and to identify the responsible genes and regulation mechanisms, we isolated, assessed, and sequenced marine microbes commonly associated with *H. echinata* (7, 8). *Shewanella* sp. strain P1-14-1 was isolated from the tissue from a feeding polyp of the marine hydroid *H. echinata*, purchased from the Marine Biological Laboratory in Woods Hole, MA. We first tested larval settlement and morphogenic activity using a larva-based assay, and then its antimicrobial activity was assessed. *Shewanella* sp. P1-14-1 showed high and reliable morphogenic activity, with a metamorphosis rate of up to 80%, and moderate antibacterial activity against Gram-positive human-pathogenic bacteria. To investigate the genetic basis of potential antimicrobial secondary metabolites (9) and the molecular details of the morphogenic activity, the genome was sequenced. Genomic DNA was extracted using the GenElute blood genomic DNA kit (Sigma-Aldrich), according to the manufacturer's protocol. Sequencing was performed at the Harvard Medical School Biopolymers Facility using Illumina TruSeq 50-bp paired-end libraries and a HiSeq 2000 flow cell (CASAVA 1.8.2; Illumina). DNA sequence data were assembled using the A5 pipeline version 20120518 (10) and analyzed for potential contaminations using Blobology (11). Genome annotation was performed using Prokka version 1.8 (12), and the G+C content was determined using QUAST version 3.0 (13). The draft genome of P1-14-1 is 4,916,416 bases in length; the G+C content

is 40.78%, with a total of 150 contigs. The largest contig assembled was 301,078 bp. The total coverage of the genome is 46-fold. Genome annotation resulted in 4,159 coding sequences (CDSs), 75 tRNAs, and 5 rRNAs. Genes associated with biofilm formation and surface attachment, including genes encoding flagella, curli, type II secretion system, type IV pili, and capsular polysaccharide (O-antigen) proteins were identified, reflecting the adaptation to successful persistence and competition on aquatic surfaces (14). The production of pili and flagella has also been connected with enhanced electrogenesis in bacteria (15, 16). Genes encoding bacteriocins and secondary metabolites (e.g., antibiotic pyrrolnitrin), were detected using antiSMASH (17) and SMIPS (18).

The draft genome sequence of *Shewanella* sp. P1-14-1 will promote the genetic analysis of the *Shewanella* genus and provide insights into secondary metabolite production and the genetic basis of the bacterial signals, which induce the settlement process of *H. echinata*.

Nucleotide sequence accession numbers. The *Shewanella* sp. P1-14-1 whole-genome shotgun (WGS) project has the project accession no. [LKTL00000000](https://ncbi.nlm.nih.gov/submit/submitseq/). The version described in this paper is the first version, with the accession number LKTL01000000.

ACKNOWLEDGMENTS

We thank the NIH for financial support to J.C. (grant GM086258), the German National Academy of Sciences Leopoldina for a postdoctoral fellowship to C.B. (grant LPDS 2011-2), and the University of Connecticut for support (J.L.K.). M.R. was supported by the graduate school Jena School for Microbial Communication (JSMC) financed by the Deutsche Forschungsgemeinschaft, and T.W. was supported by the International Leibniz Research School for Microbial and Molecular Interactions (ILRS), as part of the JSMC.

FUNDING INFORMATION

HHS | National Institutes of Health (NIH) provided funding to Jon Clardy under grant number GM086258. Deutsche Forschungsgemeinschaft (DFG) provided funding to Maja Rischer. University of Connecticut (UConn) provided funding to Jonathan Klassen. Leibniz-Gemeinschaft (Leibniz Association) provided funding to Thomas Wolf, Huijuan Guo, Ekaterina Shelest, and Christine Beemelmanns.

REFERENCES

- Ivanova EP, Sawabe T, Gorshkova NM, Svetashev VI, Mikhailov VV, Nicolau DV, Christen R. 2001. *Shewanella japonica* sp. nov. Int J Syst Evol Microbiol 51:1027–1033. <http://dx.doi.org/10.1099/00207713-51-3-1027>.
- Fredrickson JK, Romine MF, Beliaev AS, Auchtung JM, Driscoll ME, Gardner TS, Nealson KH, Osterman AL, Pinchuk G, Reed JL, Rodionov DA, Rodrigues JL, Saffarini DA, Serres MH, Spormann AM, Zhulin IB, Tiedje JM. 2008. Towards environmental systems biology of *Shewanella*. Nat Rev Microbiol 6:592–603. <http://dx.doi.org/10.1038/nrmicro1947>.
- Lovley DR. 2012. Electromicrobiology. Annu Rev Microbiol 66:391–409. <http://dx.doi.org/10.1146/annurev-micro-092611-150104>.
- Yang JL, Shen PJ, Liang X, Li YF, Bao WY, Li JL. 2013. Larval settlement and metamorphosis of the mussel *Mytilus coruscus* in response to mono-specific bacterial biofilms. Biofouling 29:247–259. <http://dx.doi.org/10.1080/08927014.2013.764412>.
- Hadfield MG. 2011. Biofilms and marine invertebrate larvae: what bacteria produce that larvae use to choose settlement sites. Annu Rev Mar Sci 3:453–470. <http://dx.doi.org/10.1146/annurev-marine-120709-142753>.
- Frank U, Leitz T, Müller WA. 2001. The hydroid *Hydractinia*: a versatile, informative cnidarian representative. Bioessays 23:963–971. <http://dx.doi.org/10.1002/bies.1137>.
- Klassen JL, Wolf T, Rischer M, Guo H, Shelest E, Clardy J, Beemelmanns C. 2015. Draft genome sequences of six *Pseudoalteromonas* strains, P1-7a, P1-9, P1-13-1a, P1-16-1b, P1-25, and P1-26, which induce larval settlement and metamorphosis in *Hydractinia echinata*. Genome Announc 3(6):e01477–15. <http://dx.doi.org/10.1128/genomeA.01477-15>.
- Klassen JL, Rischer M, Wolf T, Guo H, Shelest E, Clardy J, Beemelmanns C. 2015. Genome sequences of three *Pseudoalteromonas* strains (P1-8, P1-11, and P1-30), isolated from the marine hydroid *Hydractinia echinata*. Genome Announc 3(6):e01380–15. <http://dx.doi.org/10.1128/genomeA.01380-15>.
- Machado H, Sonnenschein EC, Melchiorson J, Gram L. 2015. Genome mining reveals unlocked bioactive potential of marine Gram-negative bacteria. BMC Genomics 16:158. <http://dx.doi.org/10.1186/s12864-015-1365-z>.
- Tritt A, Eisen JA, Facciotti MT, Darling AE. 2012. An integrated pipeline for *de novo* assembly of microbial genomes. PLoS One 7:e42304. <http://dx.doi.org/10.1371/journal.pone.0042304>.
- Kumar S, Jones M, Koutsovoulos G, Clarke M, Blaxter M. 2013. Blobology: exploring raw genome data for contaminants, symbionts and parasites using taxon-annotated GC-coverage plots. Front Genet 4:237. <http://dx.doi.org/10.3389/fgene.2013.00237>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068–2069. <http://dx.doi.org/10.1093/bioinformatics/btu153>.
- Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. Bioinformatics 29:1072–1075. <http://dx.doi.org/10.1093/bioinformatics/btt086>.
- Stevenson LH. 1977. A case for bacterial dormancy in aquatic systems. Microb Ecol 4:127–133.
- Reguera G, McCarthy KD, Mehta T, Nicoll JS, Tuominen MT, Lovley DR. 2005. Extracellular electron transfer via microbial nanowires. Nature 435:1098–1101. <http://dx.doi.org/10.1038/nature03661>.
- Gorby YA, Yanina S, McLean JS, Rosso KM, Moyles D, Dohnalkova A, Beveridge TJ, Chang IS, Kim BH, Kim BH, Kim KS, Culley DE, Reed SB, Romine MF, Saffarini DA, Hill EA, Shi L, Elias DA, Kennedy SW, Oinichuk G, Watanabe K, Ishii S, Logan B, Nealson K, Fredrickson JK, Culley DE. 2006. Electrically conductive bacterial nanowires produced by *Shewanella oneidensis* strain MR-1 and other microorganisms. Proc Natl Acad Sci USA 103:11358–11363. <http://dx.doi.org/10.1073/pnas.0604517103>.
- Weber T, Blin K, Duddela S, Krug D, Kim HU, Brucoleri R, Lee SY, Fischbach MA, Müller R, Wohlleben W, Breitling R, Takano E, Medema MH. 2015. antiSMASH 3.0—a comprehensive resource for the genome mining of biosynthetic gene clusters. Nucleic Acids Res 43:W237–W243.
- Wolf T, Shelest V, Nath N, Shelest E. 2015. CASSIS and SMIPS—promoter-based prediction of secondary metabolite gene clusters in eukaryotic genomes. Bioinformatics, in press. <http://dx.doi.org/10.1093/bioinformatics/btv713>.